REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)		
July 2014	Briefing Charts	July 2014- August 2014		
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER			
		In-House		
Novel Coordination Chemistry of Aluminus	5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)	5d. PROJECT NUMBER			
Stefan Schneider, Steve Deplazes, Yon Beauchamp	5e. TASK NUMBER			
		5f. WORK UNIT NUMBER Q0RA		
7. PERFORMING ORGANIZATION NAME(8. PERFORMING ORGANIZATION REPORT NO.			
Air Force Research Laboratory (AFMC	C)			
AFRL/RQRP				
10 E. Saturn Blvd				
Edwards AFB CA 93524-7680				
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR		10. SPONSOR/MONITOR'S ACRONYM(S)		
5 Pollux Drive.		11. SPONSOR/MONITOR'S REPORT		
Edwards AFB CA 93524-7048	NUMBER(S)			
		AFRL-RQ-ED-VG-2014-206		

12. DISTRIBUTION / AVAILABILITY STATEMENT

Distribution A: Approved for Public Release; Distribution Unlimited

13. SUPPLEMENTARY NOTES

Briefing Charts presented at ACS Fall National Meeting, San Francisco, CA, 10-14 Aug 2014. PA#14361

14. ABSTRACT

Due to its high hydrogen content, aluminum borohydride(Al(BH4)3) (ABH) makes an attractive scaffold to build upon for ionic liquids used in rocket propulsion. Due to its highly pyrophoric nature ABH poses extreme handling hazards. This reactivity can be significantly tamed through the coordination of various ligands. Previously we investigated an IL based upon the $[Al(BH_4)_4]^-$ anion. This material showed a much improved air and moisture stability compared to ABH. Here we present research on the coordination of the cyanoborohydride anion (NCBH3-) with ABH. The coordination chemistry of this anion is by far more complex than that of the simple BH4- anion and led to the discovery of, new di- and tri-anions of aluminum.

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Stefan Schneider	
a. REPORT	b. ABSTRACT	c. THIS PAGE	SAR	26	19b. TELEPHONE NO (include area code) 661-275-5759
Unclassified	Unclassified	Unclassified	57110		




Novel Coordination Chemistry of Aluminum Borohydride

ACS Fall National Meeting Aug 10 -14th 2014 S. Deplazes Edwards AFB, CA




Hydrazine – A state of the art rocket fuel





Hydrazine

Monomethylhydrazine



- Hydrazine fuel vapor toxicity can increase testing/operations costs:
 - System Handling/Fueling by certified crews in high level PPE
 - Monitoring system in field
- Vapor toxicity can limit transportation options

Ionic Liquid fuels can eliminate vapor toxicity and possess acceptable safety properties



Use of hydrazine is getting more stringent




☐ As of today, most of our in-space propulsion systems are powered by the MMH and NTO bi-propellant system, known to be hypergolic and severely toxic.
☐ In the frame of REACH (Registration, Evaluation, Authorization and Restriction of Chemicals), the European Union is enforcing the utilization of hazardous chemicals.
☐ In the mid- to long-term, the commercial use of MMH/NTO or its derivatives such as hydrazine may thus be strongly limited or even prohibited.
☐ As of June 2011 hydrazine has been added to the list of SVHC (Substance of Very High Concerns)
☐ The listing controls its use for corrosion inhibition, plating on plastics, and as a rocket fuel.



Use of Borohydride Based Salts








$AI(BH_4)_4^-$ - PROMOTES LIQUIDUS



A viscous oil crystallizing very slowly, from which neither H_2 , B_2H_6 , nor Al(BH₄)₃ could be removed even by pumping at 60°C.





Melting point depression of 75 °C.




Tetraethylammonium tetrakis(tetrahydroborato)aluminate










 $C_8N_1H_{36}AI_1B_{4,}$ 216.60 g/mol, ρ = 0.81 g/cm³ decomposition onset ~ 150°C 36.28 g/mol H in ILABH = 16.7% or 0.135 g/cm³ ~ 99% more H than LH₂/mL



Trihexyltetradecylphosphonium tetrakis(tetrahydroborato)aluminate





DISTRIBUTION A: Approved for public release; Distribution unlimited



Lack of heterocyclic BH₄ salts




- Published routes to BMIM BH₄ used IL halide in acetonitrile or CH₂Cl₂
- This work could not be reproduced and only yielded material with substantial halide content

Best results 77.5% [BH₄] halide content 22.5%

M. Bürchner, A.M.T. Erle, H. Scherer, I. Krossing Chem. Eur. J. 2012, 18, 2254.


Developed new room temperature process which yields pure materials







Anion Alteration










Distribution A: Public Release, Distribution unlimited



Cyanoborohydride coordination






Distribution A: Public Release, Distribution unlimited



X-ray crystal structure analyses as tool of characterization









Surprise! tetrakis(tetrahydroborato)aluminates










A general path to tetrakis(tetrahydroborato)aluminates?






What happened to $AI(BH_4)_2BH_3CN$?










11B NMR of reaction mixture








Maybe Chemistry is more complicated








Spiking reaction mixture with CBH









Single crystal X-ray structure analysis provided the answer



- A total of four different crystal shapes were identified under a microscope.
- Super thin plates are not a preferred crystal shape for X-ray analysis.




Anion enlarged and rotated





The real picture of the crude reaction mixture







Heat of reaction calculations




* Gas phase; all values are kcal/mol



What Happens with a Large Excess of CBH?








What Happens with a Large Excess of CBH?








Crystal Structure Shows Product





Bond	Angle (°)	Bond	Angle (°)	Bond	Length (Å)
AI1-N1-C58	178.17(16)	N1-C58-B1	178.1(2)	Al1-N2	1.9734(16)
AI1-N2-C59	174.11(15)	N2-C59-B2	178.06(18)	AI1-N3	1.9577(16)
AI1-N3-C60	172.72(15)	N3-C60-B3	179.1(2)	AI1-N6	1.9512(17)
AI1-N4-C61	175.67(16)	N4-C61-B4	178.5(2)	Al1-N4	1.9623(16)
AI1-N5-C62	174.92(16)	N5-C62-B5	179.4(2)	AI1-N5	1.9563(17)
AI1-N6-C63	177.09(17)	N6-C63-B6	177.8(3)	AI1-N1	1.9697(16)



The current reaction sequence



$$3[NCBH_{3}]^{-} + AI(BH_{4})_{3} \longrightarrow [AI(BH_{4})_{4}]^{-} + [AI(BH_{4})_{2}(NCBH_{3})_{3}]^{2-}$$

$$= \text{exc } NCBH_{3}^{-} \longrightarrow [AI(NCBH_{3})_{6}]^{3-}$$

$$2[NCBH_{3}]^{-} \longrightarrow [BH_{3}CNBH_{3}]^{-*} + [CN]^{-}$$

* Emri, J et. al., Polyhedron, 1994, 13, 2353

Summary and Conclusion

- The reactivity of aluminum borohydride is not always predictable
- Demonstrated dependence on the reaction partner and concentration
- It is challenging to characterize compound mixtures
- New species need to be isolated and incorporated into IL's to evaluate their reactivity and physical properties
- New synthetic routes to heterocyclic BH₄ salts open new possibilities



Acknowledgement





Tom Hawkins

Jeff Mills

Stefan Schneider

Capt. Andrew Beauchamp

Yonis Ahmed

Daniel Ditzhazy

Christina Franquera

ERC

Air Force Office of Scientific Research (AFOSR)



Michael Berman